



## The Smallest Gap Between Primes

---

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

May 3, 2023

# The smallest gap between primes

Frank Vega<sup>1\*</sup>

<sup>1\*</sup>Research Department, NataSquad, 10 rue de la Paix, Paris,  
75002, France.

Corresponding author(s). E-mail(s): [vega.frank@gmail.com](mailto:vega.frank@gmail.com);

## Abstract

A prime gap is the difference between two successive prime numbers. Two is the smallest possible gap between primes. A twin prime is a prime that has a prime gap of two. The twin prime conjecture states that there are infinitely many twin primes. This conjecture has been one of the great open problems in number theory for many years. In May 2013, the popular Yitang Zhang's paper was accepted by the journal *Annals of Mathematics* where it was announced that for some integer  $N$  that is less than 70 million, there are infinitely many pairs of primes that differ by  $N$ . A few months later, James Maynard gave a different proof of Yitang Zhang's theorem and showed that there are infinitely many prime gaps with size of at most 600. A collaborative effort in the Polymath Project, led by Terence Tao, reduced to the lower bound 246 just using Zhang and Maynard results as the main theoretical background. In this note, using arithmetic operations, we prove that the twin prime conjecture is true. Indeed, this is a trivial and short note very easy to check and understand which is a breakthrough result at the same time.


**Keywords:** Twin prime conjecture, Prime numbers, Prime gap

**MSC Classification:** 11A41 , 11A25

## 1 Introduction

Leonhard Euler studied the following value of the Riemann zeta function (1734).

2 *The smallest gap between primes*



**Fig. 1** Roots of  $H_2(x)$  [5]

**Proposition 1** *It is known that*[1, (1) pp. 1070]:

$$\zeta(2) = \prod_{k=1}^{\infty} \frac{p_k^2}{p_k^2 - 1} = \frac{\pi^2}{6},$$

where  $p_k$  is the  $k$ th prime number (We also use the notation  $p_n$  to denote the  $n$ th prime number).

Franz Mertens obtained some important results about the constants  $B$  and  $H$  (1874). We define  $H = \gamma - B$  such that  $B \approx 0.2614972128$  is the Meissel-Mertens constant and  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant [2, (17.) pp. 54].

**Proposition 2** *We have* [3, Lemma 2.1 (1) pp. 359]:

$$\sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \frac{1}{p_k} \right) = \gamma - B = H,$$

where  $\log$  is the natural logarithm.

For  $x \geq 2$ , the function  $u(x)$  is defined as follows [4, pp. 379]:


$$u(x) = \sum_{p_k > x} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \frac{1}{p_k} \right).$$

We use the following function:

**Definition 1** For all  $x > 1$  and  $a \geq 0$ , we define the function:

$$H_a(x) = \log\left(\frac{x}{x-1}\right) - \frac{1}{x+a} + \log\left(\frac{x^2 - \frac{\log(x)+1}{\sqrt{x}}}{x^2}\right).$$

We state the following Propositions:



**Fig. 2** Roots of  $H_4(x)$  [6]

**Proposition 3** For a sufficiently large positive value  $x$ , we have  $H_2(x) < 0$ . Certainly,  $H_2(x)$  is negative for all  $x \geq 60000$  since it is negative for  $x = 60000$ , strictly decreasing for  $x \geq 60000$  (because its derivative is lesser than 0 for  $x \geq 60000$ ) and its greatest root is between 50000 and 60000 (See Figure 1).

**Proposition 4** For a sufficiently large positive value  $x$ , we have  $H_4(x) > 0$ . Certainly,  $H_4(x)$  is positive for all  $x \geq 1.5$  since it is positive for  $x = 1.5$  and its unique root is between 1.4 and 1.5 (See Figure 2).

The following property is based on natural logarithms:

**Proposition 5** [7, pp. 1]. For  $x > -1$ :

$$\log(1+x) \leq x.$$

Putting all together yields the proof of the main theorem.

**Theorem 1** The twin prime conjecture is true.

## 2 Infinite Sums

**Lemma 1**

$$\sum_{k=1}^{\infty} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) = \log(\zeta(2)) - H.$$

## Twin prime conjecture

### 4 The smallest gap between primes

*Proof* We obtain that

$$\begin{aligned}
 \log(\zeta(2)) - H &= \log\left(\prod_{k=1}^{\infty} \frac{p_k^2}{p_k^2 - 1}\right) - H \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k^2}{p_k^2 - 1}\right) \right) - H \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k^2}{(p_k - 1) \cdot (p_k + 1)}\right) \right) - H \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) + \log\left(\frac{p_k}{p_k + 1}\right) \right) - H \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(\frac{p_k + 1}{p_k}\right) \right) - H \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_k}\right) \right) - \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \frac{1}{p_k} \right) \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_k}\right) - \log\left(\frac{p_k}{p_k - 1}\right) + \frac{1}{p_k} \right) \\
 &= \sum_{k=1}^{\infty} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right)
 \end{aligned}$$

by Propositions 1 and 2. □

#### Lemma 2

$$\sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) = \log(\zeta(2)) + \log\left(\frac{3}{2}\right).$$

*Proof* We obtain that

$$\begin{aligned}
 \log(\zeta(2)) + \log\left(\frac{3}{2}\right) &= \log(\zeta(2)) - H + H + \log\left(\frac{3}{2}\right) \\
 &= \sum_{k=1}^{\infty} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) + H + \log\left(\frac{3}{2}\right) \\
 &= \sum_{k=1}^{\infty} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) + \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \frac{1}{p_k} \right) + \log\left(\frac{3}{2}\right) \\
 &= \sum_{k=1}^{\infty} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) + \log\left(\frac{p_k}{p_k - 1}\right) - \frac{1}{p_k} \right) + \log\left(\frac{3}{2}\right) \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_k}\right) \right) + \log\left(\frac{3}{2}\right) \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right)
 \end{aligned}$$

by Lemma 1. □

### 3 Partial Sums

**Lemma 3**

$$\sum_{p_k \leq x} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) = \log\left( \prod_{p_k \leq x} \frac{p_k^2}{p_k^2 - 1} \right) - H + u(x).$$

*Proof* We obtain that

$$\begin{aligned} \log\left( \prod_{p_k \leq x} \frac{p_k^2}{p_k^2 - 1} \right) - H + u(x) &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k^2}{p_k^2 - 1} \right) \right) - H + u(x) \\ &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k^2}{(p_k - 1) \cdot (p_k + 1)} \right) \right) - H + u(x) \\ &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k}{p_k - 1} \right) + \log\left( \frac{p_k}{p_k + 1} \right) \right) - H + u(x) \\ &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \log\left( \frac{p_k + 1}{p_k} \right) \right) - H + u(x) \\ &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \log\left(1 + \frac{1}{p_k}\right) \right) - \sum_{p_k \leq x} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \frac{1}{p_k} \right) \\ &= \sum_{p_k \leq x} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \log\left(1 + \frac{1}{p_k}\right) - \log\left( \frac{p_k}{p_k - 1} \right) + \frac{1}{p_k} \right) \\ &= \sum_{p_k \leq x} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) \end{aligned}$$

by Propositions 1 and 2.  $\square$

**Lemma 4**

$$\sum_{p_k < p_n} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) = \log\left(\frac{3}{2}\right) + \log\left( \prod_{p_k \leq p_{n-1}} \frac{p_k^2}{p_k^2 - 1} \right) - \log\left(1 + \frac{1}{p_n}\right).$$

*Proof* We obtain that

$$\begin{aligned} \log\left(\frac{3}{2}\right) + \log\left( \prod_{p_k \leq p_{n-1}} \frac{p_k^2}{p_k^2 - 1} \right) - \log\left(1 + \frac{1}{p_n}\right) \\ &= \log\left(\frac{3}{2}\right) + \log\left( \prod_{p_k \leq p_{n-1}} \frac{p_k^2}{p_k^2 - 1} \right) - H + u(p_{n-1}) + H - u(p_{n-1}) - \log\left(1 + \frac{1}{p_n}\right) \\ &= \log\left(\frac{3}{2}\right) + \sum_{p_k \leq p_{n-1}} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) + H - u(p_{n-1}) - \log\left(1 + \frac{1}{p_n}\right) \\ &= \log\left(\frac{3}{2}\right) + \sum_{p_k \leq p_{n-1}} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) \right) + \sum_{p_k \leq p_{n-1}} \left( \log\left( \frac{p_k}{p_k - 1} \right) - \frac{1}{p_k} \right) - \log\left(1 + \frac{1}{p_n}\right) \end{aligned}$$

## Twin prime conjecture

### 6 The smallest gap between primes

$$\begin{aligned}
 &= \log\left(\frac{3}{2}\right) + \sum_{p_k \leq p_{n-1}} \left( \frac{1}{p_k} - \log\left(1 + \frac{1}{p_k}\right) + \log\left(\frac{p_k}{p_k-1}\right) - \frac{1}{p_k} \right) - \log\left(1 + \frac{1}{p_n}\right) \\
 &= \log\left(\frac{3}{2}\right) + \sum_{p_k \leq p_{n-1}} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_k}\right) \right) - \log\left(1 + \frac{1}{p_n}\right) \\
 &= \sum_{p_k < p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right)
 \end{aligned}$$

by Lemma 3. □

## 4 Main Insight

**Lemma 5**

$$\sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) = \log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2-1}\right).$$

*Proof* We obtain that

$$\begin{aligned}
 &\sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) \\
 &= \sum_{k=1}^{\infty} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) - \sum_{p_k < p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) \\
 &= \log(\zeta(2)) + \log\left(\frac{3}{2}\right) - \log\left(\frac{3}{2}\right) - \log\left(\prod_{p_k \leq p_{n-1}} \frac{p_k^2}{p_k^2-1}\right) + \log\left(1 + \frac{1}{p_n}\right) \\
 &= \log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2-1}\right)
 \end{aligned}$$

by Lemmas 2 and 4. □

## 5 Proof of Theorem 1

*Proof* Suppose that the twin prime conjecture is false. Then, there would exist a sufficiently large prime number  $p_n$  such that for all prime gaps starting from  $p_n$ , this implies that they are greater than or equal to 4. First, we need to prove that

$$\log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2-1}\right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}{p_k^2}\right) \leq 0$$

when  $p_n$  is large enough. That is the same as

$$\log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2-1}\right) \leq \sum_{p_k \geq p_n} \log\left(\frac{p_k^2}{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}\right).$$

That is equivalent to

$$\frac{p_n + 1}{p_n} \cdot \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2-1} \leq \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}.$$

So,

$$\frac{p_n + 1}{p_n} \cdot \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 1} \leq \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - \left( \frac{\log(p_k) + 1}{\sqrt{p_k}} \right)^{\frac{\log(p_k)}{\log(p_k)}}}$$

which is

$$\frac{p_n + 1}{p_n} \cdot \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 1} \leq \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - \frac{\log(p_k) + 1}{\log(p_k) \sqrt{e}}}$$

since  $x^{\frac{1}{\log x}} = e$  for  $x > -1$ . Hence, it is enough to show that

$$\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 2.71828} \leq \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - \frac{\log(p_k) + 1}{\log(p_k) \sqrt{e}}}$$

for a sufficiently large prime number  $p_n$ . Indeed, we see that

$$(p_n + 1) \cdot \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 1} \leq p_n \cdot \prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 2.71828}$$

trivially holds for large enough  $p_n$  since

$$\frac{p_n \cdot (p_n - 1)}{p_n^2 - 1} \geq \prod_{p_k \geq p_n} \frac{p_k^2 - 2.71828}{p_k^2 - 1}.$$

We deduce that

$$\frac{p_n^4 - 2.71828}{p_n^4 - 1} \geq \prod_{p_k \geq p_n} \frac{p_k^2 - 2.71828}{p_k^2 - 1}$$

since there is always a  $k'$  such that  $p_n < p_{k'} < p_n^2$  and

$$\frac{p_n^4 - 2.71828}{p_n^4 - 1} > \frac{p_{k'}^2 - 2.71828}{p_{k'}^2 - 1} > \prod_{p_k \geq p_n} \frac{p_k^2 - 2.71828}{p_k^2 - 1}$$

because of

$$\frac{p_k^2 - 2.71828}{p_k^2 - 1} < 0$$

for every  $k$ . Thus, we only need to prove that

$$\frac{p_n \cdot (p_n - 1)}{p_n^2 - 1} \geq \frac{p_n^4 - 2.71828}{p_n^4 - 1}$$

that is simply

$$p_n \cdot (p_n - 1) \cdot (p_n^2 + 1) \geq p_n^4 - 2.71828$$

which is trivially true for a sufficiently large prime number  $p_n$ . Consequently, we can assure that

$$\log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 1}\right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k) + 1}{\sqrt{p_k}}}{p_k^2}\right) \leq 0$$

whenever  $p_n$  is large enough. In this way, we have

$$\sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k - 1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k) + 1}{\sqrt{p_k}}}{p_k^2}\right) \leq 0$$



8 *The smallest gap between primes*

by Lemma 5. We verify that

$$\begin{aligned} & \sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \log\left(1 + \frac{1}{p_{k+1}}\right) \right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}{p_k^2}\right) \\ & \geq \sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \frac{1}{p_{k+1}} \right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}{p_k^2}\right) \end{aligned}$$

since  $-\log\left(1 + \frac{1}{p_{k+1}}\right) \geq -\frac{1}{p_{k+1}}$  by Proposition 5. Under our assumption, we notice that

$$\sum_{p_k \geq p_n} \left( \log\left(\frac{p_k}{p_k-1}\right) - \frac{1}{p_{k+1}} \right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}{p_k^2}\right) \geq \sum_{p_k \geq p_n} H_4(p_k)$$

since  $-\frac{1}{p_{k+1}} \geq -\frac{1}{p_{k+4}}$ . However, we know that

$$\sum_{p_k \geq p_n} H_4(p_k) > 0$$

due to Proposition 4. Hence, the inequality

$$\log\left(1 + \frac{1}{p_n}\right) + \log\left(\prod_{p_k \geq p_n} \frac{p_k^2}{p_k^2 - 1}\right) + \sum_{p_k \geq p_n} \log\left(\frac{p_k^2 - \frac{\log(p_k)+1}{\sqrt{p_k}}}{p_k^2}\right) \leq 0$$

would not hold by transitivity. For that reason, we obtain a contradiction under the supposition that the twin prime conjecture is false. By *reductio ad absurdum*, we prove that the twin prime conjecture is true.  $\square$

## References

- [1] R. Ayoub, Euler and the zeta function. *The American Mathematical Monthly* **81**(10), 1067–1086 (1974). <https://doi.org/10.2307/2319041>
- [2] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie. *J. reine angew. Math.* **1874**(78), 46–62 (1874). <https://doi.org/10.1515/crll.1874.78.46>
- [3] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin’s criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux* **19**(2), 357–372 (2007). <https://doi.org/10.5802/jtnb.591>
- [4] J.L. Nicolas, Petites valeurs de la fonction d’Euler. *Journal of number theory* **17**(3), 375–388 (1983). [https://doi.org/10.1016/0022-314X\(83\)90055-0](https://doi.org/10.1016/0022-314X(83)90055-0)
- [5] Equation Solver - Wolfram Alpha. Roots for the function H in the value of  $a = 2$ . <https://www.wolframalpha.com/input?i2d=true&i=log%5C%2840%29Divide%5BX%2C%5C%2840%29X+-+1%5C%2841%29%5D%5C%2841%29+-+Divide%5B1%2C%5C%2840%29X+%2B+2%5C%2841%29%5D+%2B+log%5C%2840%29Divide%5B%5C%2840%29Power%5BX%2C%5D+-+Power%5BX%2C%5C%2840%29Divide%5B1%2C%5C%2840%29log%5C%2840%29X%5C%2841%29+%2B+1>

